


Cosmic Microwave Background *and* CMB-S4

CMB-S4: a coherent project building on CMB stage II & III projects including participation by:

- the ACT, BICEP/KECK, SPT and Polarbear CMB teams;
- Argonne, FNAL, LBNL and SLAC national labs;
- and the HEP community.

John Carlstrom
for the CF5 Inflation and Neutrino topical groups
and the CMB-S4 collaboration

Early universe as a HEP lab




Inflation?

Universe expands by $>e^{60}$
solving smoothness problem,
flatness and more..

What drove inflation?

What is the energy scale of inflation?


- spectral index of fluctuations, n_s
- constrain tensor to scalar fluctuations
- inflationary gravitational wave B-mode polarization
- non-Gaussianity?





Neutrinos?

N_{eff}

Effective number of relativistic species
'dark radiation' impacts intrinsic CMB power spectrum





Neutrinos?



$$\Sigma m_\nu$$

Sum of the neutrino masses
impacts growth of large scale structure,
i.e., the matter power spectrum
Probed by CMB lensing

Neutrinos?




$$\Sigma m_\nu > 0$$

Sum of the neutrino masses
impacts growth of large scale structure,
i.e., the matter power spectrum
Probed by CMB lensing


WMAP

$\sim 70 \text{ deg}^2$



Planck 143 GHz

$\sim 70 \text{ deg}^2$




Ground based 150 GHz (SPTpol)

~70 deg²

13x higher resolution and 60x deeper than WMAP


7x higher resolution and 9x deeper than Planck




Ground based 150 GHz (SPTpol)

$\sim 70 \text{ deg}^2$

**CMB-S4 will be 5x deeper and cover
50x more sky than SPTpol survey**




Primary CMB anisotropy - 9 harmonics



Inflation checks: Geometrically flat universe; Superhorizon features; Acoustic peaks/adiabatic fluctuations; Departure from scale invariance.


Constraining inflationary models joint r and n_s limits



Spectral Index of primordial fluctuations, n_s ,
where $\Delta_R^2(k) = \Delta_R^2(k_0) \left(\frac{k}{k_0} \right)^{n_s-1}$

Inflation evidence
 $n_s \neq 1$ at over 5σ

Constraining inflationary models joint r and n_s limits




Spectral Index of primordial fluctuations, n_s ,
where $\Delta_R^2(k) = \Delta_R^2(k_0) \left(\frac{k}{k_0} \right)^{n_s-1}$


Inflation evidence
 $n_s \neq 1$ at over 5σ

Primary CMB anisotropy - 9 harmonics

Improves precision of sound horizon, θ_s ,
& provides larger lever arm




And most importantly provides determination of the damping scale, θ_d




Note $\frac{r_d}{r_s} = \frac{\theta_d}{\theta_s} \propto H^{0.5}$ so ratio is sensitive to energy density.

Constraining model extensions: joint N_{eff} and Σm_ν constraints




N_{eff} is the effective number of relativistic species.
For standard 3 neutrinos $N_{\text{eff}} = 3.046$.
It measures the extra energy relative to the photons.

Constraining model extensions: joint N_{eff} and Σm_ν constraints



N_{eff} is the effective number of relativistic species.
For standard 3 neutrinos $N_{\text{eff}} = 3.046$.
It measures the extra energy relative to the photons.

Constraining model extensions: joint N_{eff} and Σm_ν constraints



N_{eff} is the effective number of relativistic species.
For standard 3 neutrinos $N_{\text{eff}} = 3.046$.
It measures the extra energy relative to the photons.


CMB lensing

Large-Scale Structure Lenses the CMB


- RMS deflection of $\sim 2.5'$
- Lensing efficiency peaks at $z \sim 2$
- Coherent on \sim degree (~ 300 Mpc) scales

Lensing of the CMB

$17^\circ \times 17^\circ$



lensing potential




unlensed cmb


from Alex van Engelen

Lensing of the CMB

$17^\circ \times 17^\circ$




lensing potential




lensed cmb

from Alex van Engelen

We can take a CMB map




and construct the CMB Lensing map
reconstruction of the mass projected
along the line of sight to the CMB.




Lensing convergence map smoothed to 1 deg resolution

and construct the CMB Lensing map
reconstruction of the mass projected
along the line of sight to the CMB.




Correlation of matter traced by CMB lensing (contours) and distribution of high z galaxies (grayscale; Herschel 500 μm) [arXiv:1112.5435]



Lensing convergence map smoothed to 1 deg resolution

CMB lensing power spectrum

Planck XVII 2013




Sensitive to the neutrino masses
 $\Sigma m_\nu = 0.1 \text{ eV} \rightarrow 5\% \text{ amplitude of spectrum}$

Polarization gives additional lensing sensitivity and is a cleaner probe.

CMB lensing power spectrum

Planck XVII 2013



Sensitive to the neutrino masses
 $\Sigma m_\nu = 0.1 \text{ eV} \rightarrow 5\% \text{ amplitude of spectrum}$


Polarization gives additional lensing sensitivity and is a cleaner probe.

CMB lensing and optical surveys

CMB lensing complements large optical surveys such as DES, eBOSS, LSST, DESI, Euclid, WFIRST, etc.


The combination leads to better shear-bias calibration and more robust constraints on Dark Energy and the properties of neutrinos.

→Critical for CMB-S4 sky coverage to overlap optical surveys.




From “Can CMB Lensing Help Cosmic Shear Surveys?” Das, Errard, and Spergel, 2013


CMB polarization: ***the next frontier for lensing & inflation***




CMB polarization: ***the next frontier for lensing & inflation***




CMB polarization: ***the next frontier for lensing & inflation***



CMB polarization: ***the next frontier for lensing & inflation***



CMB polarization: ***the next frontier for lensing & inflation***



CMB polarization: ***the next frontier for lensing & inflation***









Figure from CF5 inflation doc: note expanded scale with $0.001 < r < 0.01$

Status of B-mode experiments

Barkats et al., arXiv:1310.1422



SPTpol Detection of lensing B-modes



CMB timeline

- **2009**: $r < 0.7$ (BICEP) Chiang et al, 0906.1181

- **2013**: Stage II experiments detect lensing B-modes
- **2014**: $r \lesssim 0.1$ from Inflationary B-modes (BICEP 2) ?
- **2013-2016**: Stage II experiments
 $\sigma(r) \sim 0.03$, $\sigma(N_{\text{eff}}) \sim 0.1$, $\sigma(\Sigma m_\nu) \sim 0.1 \text{ eV}$
- **2016-2020**: Stage III experiments
 $\sigma(r) \sim 0.01$, $\sigma(N_{\text{eff}}) \sim 0.06$, $\sigma(\Sigma m_\nu) \sim 0.06 \text{ eV}$;

- **2020-2025**: Stage IV experiment, **CMB-S4**
 $\sigma(r) = 0.001$, $\sigma(N_{\text{eff}}) = 0.020$, $\sigma(\Sigma m_\nu) = 16 \text{ meV}$

CMB timeline

- **2009**: $r < 0.7$ (BICEP) Chiang et al, 0906.1181


- **2013**: Stage II experiments detect lensing B-modes
- **2014**: $r \lesssim 0.1$ from Inflationary B-modes (BICEP 2) ?
- **2013-2016**: Stage II experiments
 $\sigma(r) \sim 0.03$, $\sigma(N_{\text{eff}}) \sim 0.1$, $\sigma(\Sigma m_\nu) \sim 0.1 \text{ eV}$
- **2016-2020**: Stage III experiments
 $\sigma(r) \sim 0.01$, $\sigma(N_{\text{eff}}) \sim 0.06$, $\sigma(\Sigma m_\nu) \sim 0.06 \text{ eV}$;

- **2020-2025**: Stage IV experiment, **CMB-S4**
 $\sigma(r) = 0.001$, $\sigma(N_{\text{eff}}) = 0.020$, $\sigma(\Sigma m_\nu) = 16 \text{ meV}$

On an ambitious path forward and producing a steady flow of scientific results

CMB-S4 What it will deliver

Inflation projection for CMB-S4




— 3.5 $\mu\text{K}\cdot\sqrt{\text{S}}$ 8'FWHM 10%FG
— 3.5 $\mu\text{K}\cdot\sqrt{\text{S}}$ 1'FWHM 5%FG
— 1.1 $\mu\text{K}\cdot\sqrt{\text{S}}$ 1'FWHM 10%FG

— 3.5 $\mu\text{K}\cdot\sqrt{\text{S}}$ 1'FWHM 10%FG
- - 1.1 $\mu\text{K}\cdot\sqrt{\text{S}}$ 8'FWHM 5%FG
- - 1.1 $\mu\text{K}\cdot\sqrt{\text{S}}$ 1'FWHM 5%FG

CMB-S4 What it will deliver

Joint projections $N_{\text{eff}} - \Sigma m_\nu$



$\sigma(\Sigma m_\nu) = 16 \text{ meV}$
with two probes!

$\sigma(N_{\text{eff}}) = 0.020$
unique to CMB

CMB-S4 What it will take

- **CMB-S4 Survey:**

- Maximum return on Inflation, Neutrino, and Dark Energy science requires an optimized survey which includes a range of resolution and sky coverage from deep to wide.

- **Sensitivity of ~ 1 μ K-arcmin over half the sky**

- **Experimental Configuration:**

- 200,000+ detectors on multiple platforms
 - spanning 40 - 240 GHz for foreground removal
 - $\lesssim 3$ arcmin resolution required for CMB lensing & neutrino science,

- (higher resolution leads to amazing and complementary dark energy constraints and gravity tests on large scales via the SZ effect)*

See Snowmass planning document [arxiv:1309.5383](https://arxiv.org/abs/1309.5383)

CMB-S4: How to do it

- Build on extensive experience from earlier generation CMB experience
 - People
 - Technology
 - Systematic Error Control
 - Analysis
- And increase throughput by over an order of magnitude

CMB-S4: How to do it

- Build on extensive experience from earlier generation CMB experience
 - People
 - Technology
 - Systematic Error Control
 - Analysis
- And increase throughput by over an order of magnitude

Technical challenge:

is the scaling of the CMB detector arrays.

Sociological evolution:


the highly competitive CMB groups are working together.

CMB-S4: How to do it


- Exploit superb, established sites at Atacama Chile and South Pole
 - proven high and dry sites for sensitive CMB measurements
 - provides the required access to $> 50\%$ of the sky, including coverage of the optical survey fields

CMB S4 Large Area Survey Region

(overlap with LSST, MS-DESI, etc)



Build on investment by NSF in Chilean mid-Latitude Facilities and CMB experiments



5200 meter (17,000 ft) site developed by the ACT team since 1998 provides access to over 50% of the sky

- The Atacama Cosmology Telescope
 - 6 meter aperture (1.4 arcmin at 150 GHz)
- Polarbear Telescope
 - 3.5 meter (3.5 arcmin at 150 GHz)



\$15M+ in Telescopes and Logistics


- Power, internet, workspace, roads
- Machine shop and supplies
- Low altitude control room and housing
- Ties to Chilean contractors and suppliers
- Legal presence in Chile
- Established positive working relationship with CONICYT (Chilean NSF)

Build on investment by NSF in South Pole Facilities and CMB experiments



- Major NSF research station (not shown) with excellent logistical support
- CMB measurements since the 1980s;
Martin A. Pomerantz observatory established in 1994.
- Exceptionally low atmospheric noise (sky-noise) due to dry and stable atmosphere.
- Access to ~4000 square degrees of low foreground sky (10%), which is observable year-round, 24 hrs/day

CMB-S4: How to do it





2012: SPTpol Stage II
1600 detectors (ANL/NIST)

ANL, LBNL, SLAC, Polarbear and SPT teams working on Stage II to Stage III detector advance based on UCB 3-band, dual polarization pixel;

- ▶ *Optimized with background limited noise and high throughput*
- ▶ *Uniform properties over 150-mm diameter wafers*
- ▶ *Consistent fabrication from batch-to-batch*

UC Berkeley prototype pixel




2016: SPT-3G Stage III 4x larger area
15,234 detectors at $T = 250\text{mK}$

CMB-S4: How to do it

Stage-3


~15,200 detectors



Stage-4

2020+: CMB-S4

200,000+ detectors
multiple telescopes



Detector sensitivity has been limited by photon “shot” noise for last ~15 years; further improvements are made only by making ***more detectors***.

CMB-S4: How to do it

→ CMB-S4 requirements exceed capabilities of the traditional University-based CMB groups

- Increased production scope and reliability
 - 200,000+ detectors requires production of approximately 150 silicon 6" detector arrays
- Multiplexed TES Readout
- Large Cryogenic Optics
- Computing Infrastructure and Analysis tools
 - ~10,000 x *Planck* data size (~ 6 TB/day)
- Project Organization/Management

→ requires DOE National lab and HEP community working with the University-based CMB groups



- **Investment in robust, large scale detector fabrication.**
- **Involvement with SPTpol Stage II experiment (provided 90 GHz channel).**
- **Involvement in SPT-3G Stage III, providing detectors.**




- **Investment in multiplexer readout.**
- **CMB heritage and connections with UCB detector development.**
- **High performance computing/ massively parallel data analysis.**
- **Involvement in Polarbear and SPT (Stage II & III).**



- **Detector testing, SiDet for module assembly, and radiometer cryostat design, testing and integration.**
- **Experience with QUIET detector module testing and assembly.**



- **Investment in developing large aperture cryogenic optics.**
- **Investing in robust, large scale detector fabrication**
- **Investment in SQuIDs.**
- **Involvement in BICEP / KECK, SPT and ACT Stage III, providing detectors.**



NOTIONAL BUDGET		FY15	FY16	FY17	FY18	FY19	FY20	
DOE ANL/LBNL/SLAC Detector	project capital \$*	0.5M	1M	1.5M	1.5M	1.5M	1.5M	
	FTE†	5	7	10	14	16	16	
DOE (LBNL/NERSC) Computing	project capital \$	-	-	-	-	-	0.5M	
	FTE	-	1	2	2	3	3	
DOE Receiver test facilities, hardware & electronics	project capital \$	-	3M	4.5M	7.5M	7.5M	7.5M	
	FTE	4	8	10	12	12	12	
NSF University CMB Dev, Test, Ops and Analysis	Σ \$	(7M) current	7M	7M	7M	8M	9M	
new telescopes (NSF) site/deploy (NSF/DOE)	project capital \$	-	2M	7M	7M	7M	7M	
NSF & DOE new telescope operations	\$	-	0.5M	2M	3M	4M	5M	
DOE Lab & Univ Analysis (converts)	FTE	6	10	16	24	30	30	
*2013 dollars †DOE Particle Physicist FTE								

NOTIONAL BUDGET		FY15	FY16	FY17	FY18	FY19	FY20	Capital \$+FTEyr
DOE ANL/LBNL/SLAC Detector	project capital \$	0.5M	1M	1.5M	1.5M	1.5M	1.5M	7.5M
	FTE	5	7	10	14	16	16	68 yr
DOE (LBNL/NERSC) Computing	project capital \$	-	-	-	-	-	0.5M	0.5M
	FTE	-	1	2	2	3	3	11 yr
DOE Receiver test facilities, hardware & electronics	project capital \$	-	3M	4.5M	7.5M	7.5M	7.5M	31M
	FTE	4	8	10	12	12	12	58 yr
NSF University CMB Dev, Test, Ops and Analysis	Σ \$	(7M) current	7M	7M	7M	8M	9M	
†new telescopes (NSF) site/deploy (NSF/DOE)	project capital \$	-	2M	7M	7M	7M	7M	30M
NSF & DOE new telescope operations	\$	-	0.5M	2M	3M	4M	5M	
DOE Lab & Univ Analysis (converts)	FTE	6	10	16	24	30	30	
†Roughly 2:1 ratio of cost of telescopes to costs of site prep and deployment		Total Project Capital: \$69M and 137 FTE • yr (not including 25% contingency)						

NOTIONAL BUDGET		FY15	FY16	FY17	FY18	FY19	FY20	Ops/yr Analysis/yr
DOE ANL/LBNL/SLAC Detector	project capital \$	0.5M	1M	1.5M	1.5M	1.5M	1.5M	
	FTE	5	7	10	14	16	16	
DOE (LBNL/NERSC) Computing	project capital \$	-	-	-	-	-	0.5M	0.5M/ 3yr
	FTE	-	1	2	2	3	3	4 FTE
DOE Receiver test facilities, hardware & electronics	project capital \$	-	3M	4.5M	7.5M	7.5M	7.5M	
	FTE	4	8	10	12	12	12	
NSF University CMB Dev, Test, Ops and Analysis	Σ \$	(7M) current	7M	7M	7M	8M	9M	9M
new telescopes (NSF) site/deploy (NSF/DOE)	project capital \$	-	2M	7M	7M	7M	7M	
NSF & DOE new telescope operations	\$	-	0.5M	2M	3M	4M	5M	5M
DOE Lab & Univ Analysis (converts)	FTE	6	10	16	24	30	30	30+FTE
		2020+ yearly ops and analysis: \$14.2M and 34+ FTE						

International competition / partners


- There is no competition at the scale of CMB-S4
 - European L-class mission (PRISM) was turned down
 - No NASA mission expected on this time scale
- International partners
 - We envision CMB-S4 as primarily a U.S. project
 - Current international partners contributing to the CMB teams, e.g.,
 - Cardiff with all
 - KEK, Japan with Polarbear
 - McGill U with Polarbear and SPT
 - CITA, Oxford, UBC with ACT
 - Chile is critical partner; ACT works with CONICYT (Chilean NSF)
 - plus many other international participants
 - We expect a lot of international interest if CMB-S4 goes forward.

What we hope P5 will endorse

1. CMB uniquely addresses fundamental and exciting HEP science.
2. DOE-HEP has critical role in current and future CMB experiments.
3. Continued NSF and DOE funding of the CMB groups is critical to advancing CMB science.
 - It is essential to include the expertise from established university CMB groups.
 - Best and most economical path to CMB-S4 is to build on existing CMB experiments & telescopes.
4. CMB-S4 technology is identified and significantly mature to push for large scale integration; CMB-S4 detector development could and should start ASAP.
5. CMB-S4 should exploit infrastructure investments in robust, large scale micro-fabrication at ANL and SLAC, and in detector development at LBNL. Two production facilities will be needed.
6. CMB-S4 program could be ready for project CD0 in 2015, with full deployment in 2020 and measurements continuing through 2024.
7. The CMB program and CMB-S4 is expected to continue to produce a steady flow of scientific results and new discoveries en route to achieving its primary goals.


Extra slides

Experimental Evolution



Combined Neutrino mass constraints


Future Cosmology
 $\sigma(\Sigma m_\nu) = 16 \text{ meV}$



“use cosmology to tighten the noose” Boris Kayser

CMB Sunyaev-Zel'dovich Cluster Survey

Cluster Mass vs Redshift from CMB SZ measurements




CMB measurements detect clusters through the “shadows” they make in the CMB, the Sunyaev-Zel'dovich (SZ) effect:

SPT-SZ/pol: $N_{\text{clust}} \sim 1,000$
SPT-3G: $N_{\text{clust}} \sim 10,000$
CMB-S4: $N_{\text{clust}} \sim 100,000+$


CMB lensing measured from individual clusters, can directly calibrate cluster mass:

SPT-3G: $\sigma(M) \sim 3\%$
CMB-S4: $\sigma(M) < \sim 0.1\%$

CMB-S4 Lensing Sensitivity Σm_ν



setting limit to tensor perturbations i.e., primordial gravitational waves



$$r \equiv \frac{\text{Tensor (gravitational) perturbation amplitude}}{\text{Scalar (density) perturbation amplitude}} \quad V^{1/4} = 1.06 \times 10^{16} \text{ GeV} \left(\frac{r}{0.01} \right)^{1/4}$$